

## AN IMPROVED ALGORITHM FOR OPTIMIZING PRODUCT MIX IN ASSIGNMENT PROBLEMS USING UNCERTAINTY PARAMETERS

P. Mathiyalagan<sup>1</sup> & R. Nagarajan<sup>2</sup>

1.Professor, Department of Mechanical Engineering, J.J College of Engineering &Technology, Tiruchirappalli-620009

2.Professor, Department of Mathematics , J.J College of Engineering &Technology, Tiruchirappalli-620009

Corresponding Author: [rajenagarajan1970@gmail.com](mailto:rajenagarajan1970@gmail.com).

ORCID ID: <https://orcid.org/0000-0002-7333-8710>

**Abstract :** Inventory models in which the demand rate on the inventory level are based on the common real life observation that greater product availability tends to stimulate more sales. Theory of constraints (TOC) is a production planning philosophy that tries to improve the throughput of the system management of inventory levels. Due to the existing of inventory levels in a production system the demands of all products can not be fully met. So one of the most important decisions made in production systems is product mix problem. Although many algorithms have been developed in the fields using the concept of theory of constraints. This paper benefits from a variety of advantages. In order to consider the importance of all inventory levels, group decision making approach is applied and the optimal product mix is reached. In the algorithm presented in this paper, each inventory level is considered as a decision maker. The new algorithm benefits from the concept of fuzzy group decision making and optimizes the product mix problem in inventory environment where all parameters are fuzzy values.

**Keywords :**Fuzzy group decision making, product mix optimization, multi-stage decision making, theory of constraints, inventory level cost, Triangular fuzzy number.

AMS Subject classification :

**Section-1 Introduction :** Theory of constraints (TOC) which has been first introduced in the Goal [4] is a production planning philosophy that aims to improve the system through put by efficient use of inventory levels. In this paper product mix optimization is

considered as a decision making problem. Regarding this analogy decision making criteria should be first defined [5]. Two important criteria are throughput and the later delivery cost. Later delivery cost is the most of mission one unit of each product. Assuming each inventory level as a decision maker [6], product mix optimization is a group decision making problem. In all previous researchers all parameters (such as processing time, demand etc) are assumed as crisp values. In this paper, a new algorithm is developed to optimize the product mix problem with all inputs are fuzzy values and Borda methods is used in group decision making process as ordinal techniques are preferred to cardinal ones [9].

### Section – 2 Algorithm.

The following notations are used in the new algorithm.

$t_{ij}$  = processing time of product  $i$  on resource  $j$ .

$D_i$  = Demand of product  $i$ .

$Sp_i$  = selling price of product  $i$ .

$Rm_i$  = Raw material cost of product  $i$ .

$Ac_j$  Available capacity of resource  $j$ .

$Rc_j$  = Required capacity of resource  $j$ .

$n$  = number of products.

$m$  = number of inventory levels.

In this paper, all parameters are considered triangular fuzzy numbers and are shown as  $(x,y,z)$  where  $x < y < z$ ,  $\mu_y = 1$  and  $\mu_x = \mu_z = 0$ . so let define  $t_{ij}$ ,  $D_i$ ,  $Sp_i$ ,  $Rm_i$ , and  $Ac_j$  as follows.

$$t_{ij} = (L_{ij}, M_{ij}, U_{ij}), \quad D_i = (L_i', M_i', U_i') \quad Sp_i = (A_i, B_i, C_i)$$

$$Rm_i = (A_i', B_i', C_i'), \quad Ac = (\alpha_i, \beta_i, \chi_i)$$

**Step – 1 :** Identify the system of inventory levels. As  $t_{ij}$  and  $D_i$  are positive fuzzy numbers, the required capacity of resource  $j$  is calculated as follows.

$$Rc_j = \sum_{i=1}^n t_{ij} D_i = \left( \sum_{i=1}^n L_{ij} L_i', \sum_{i=1}^n M_{ij} M_i', \sum_{i=1}^n U_{ij} U_i' \right) \dots\dots\dots (1)$$

For simplicity  $Rc_j$  is shown as  $(a_j, b_j, c_j)$ . In order to determine whether  $j$  is an inventory level.,  $Rc_j$  and  $Ac_j$  are compared using fuzzy ranking techniques. Due to the efficiency of ranking of ranking methods based on left and right scores, the method by chen is applied (chen and Hwang 1992). In this method, right and left scores of a fuzzy number refer to its intersection with the fuzzy max and the fuzzy min respectively. The fuzzy max and fuzzy min are defined as follows,

$$\mu_{\max}(x) = \frac{x - x_{\min}}{x_{\max} - x_{\min}} \quad \text{and} \quad \mu_{\min}(x) = \frac{x_{\max} - x}{x_{\max} - x_{\min}} \quad \dots\dots\dots (2)$$

where  $x_{\max}$  is  $\max(b_i, \chi_i)$  and  $x_{\min}$  ( $a_i, \alpha_i$ ). As  $Rc_i$  and  $Ac_i$  are triangular fuzzy numbers, they are convex, continuous and normal. So their right scores may be determined by taking the intersection of their non-increasing part and  $\mu_{\max}(x)$ . similarly their scores are determined by taking the intersection of their non-decreasing part and  $\mu_{\min}(x)$ . As higher right score  $\mu_R(x)$  and lower left score indicate large fuzzy number, the total score of  $Rc$  and  $Ac$  are defined as follows.

$$\mu_{\text{Total}}(Rc_j) = \frac{\mu_{\text{Right}}(Rc_j) + 1 - \mu_{\text{Left}}(Rc_j)}{2} \dots\dots\dots (3)$$

$$\mu_{\text{Total}}(Ac_j) = \frac{\mu_{\text{Right}}(Ac_j) + 1 - \mu_{\text{Left}}(Ac_j)}{2} \dots\dots\dots (4)$$

If  $\mu_{\text{Total}}(Rc_j) > \mu_{\text{Total}}(Ac_j)$ , then  $j$  is an inventory level.

**Step – 2 :** Form decision matrices. Throughput ( $X_{ij}$ ) is the first criterion considered in optimizing product mix. It is calculated as  $X_{ik} = Cm_i / t_{ij} \dots\dots\dots (5)$  Where  $Cm_i$  is determined by the difference of the selling price and raw material cost of the product  $i$ .

$C_{mi} = (A_i - C_i', B_i - B_i', C_i - A_i')$  ..... (6) As  $C_{mi}$  and  $t_{ij}$  are triangular fuzzy numbers,  $X_{ik}$  is calculated as follows.

$$X_{ik} = (A_i / U_{ij}, B_i / M_{ij}, C_i / L_{ij},) \dots\dots\dots (7)$$

The other criteria is late delivery cost (Rashidi Komijan and Sadjadi 2005) . Although it is a crisp value in most cases, it is considered as  $LDC_i (p_i, q_i, r_i)$ . Decision matrix of decision make K can be set as follows.

$$DM_k = \begin{pmatrix} Z_1 & X_{1k} & LDC_1 \\ \cdot & \cdot & \cdot \\ Z_i & X_{ik} & LDC_i \\ \cdot & \cdot & \cdot \\ Z_k & X_{nk} & LDC_n \end{pmatrix} \dots\dots\dots (9)$$

Where  $r_{ijk}$  is the rank of alternative I assigned by decision maker K given the criterion j.

**Step – 4:** For each criterion, set an agreed matrix that shows the ranks assigned to the alternatives by decision makers.

$$R_j' = \begin{matrix} Z_1 \\ Z_i \\ Z_n \end{matrix} \begin{pmatrix} 1 \dots\dots\dots k \dots\dots\dots m \\ r_{1j1} \dots\dots r_{1jk} \dots\dots\dots r_{1jm} \\ r_{ij1} \dots\dots r_{ijk} \dots\dots\dots r_{ijm} \\ r_{nj1} \dots\dots r_{njk} \dots\dots\dots r_{njm} \end{pmatrix} \dots\dots\dots (10)$$

**Step – 5;** Form Borda score matrices.

$$B_j' = \begin{matrix} Z_1 \\ Z_i \\ Z_n \end{matrix} \begin{pmatrix} 1 \dots\dots\dots k \dots\dots\dots m \\ b_{1j1} \dots\dots b_{1jk} \dots\dots\dots b_{1jm} \\ b_{ij1} \dots\dots b_{ijk} \dots\dots\dots b_{ijm} \\ b_{nj1} \dots\dots b_{njk} \dots\dots\dots b_{njm} \end{pmatrix} \dots\dots\dots (11)$$

Where  $b_{ijk} = n - r_{ijk}$

**Step – 6;** Set score matrices by summing the values of each row.

$$SM_j = \begin{matrix} Z_1 \\ \cdot \\ Z_i \\ \cdot \\ Z_n \end{matrix} \begin{pmatrix} S_{1j} \\ \cdot \\ S_{ij} \\ \cdot \\ S_{nj} \end{pmatrix} \dots\dots\dots (12)$$

Where  $S_{ij} = \sum_{K=1}^n b_{ijk}$

**Step – 7 :** Set the agreement matrix. Firstly, values of the score matrices should be ranked. Then the agreement matrix is set by aggregating these ranks.

$$R_G = \begin{matrix} Z_1 \\ \cdot \\ Z_i \\ \cdot \\ Z_n \end{matrix} \begin{pmatrix} G_{11} & G_{12} \\ \cdot & \cdot \\ G_{i1} & G_{i2} \\ \cdot & \cdot \\ G_{n1} & G_{n2} \end{pmatrix} \dots\dots\dots (13)$$

Where  $G_{i,1}$  and  $G_{i,2}$  are the agreement ranks of alternative ‘i’ given through put and late delivery cost respectively.

**Step – 8 :** Set the collective weighted agreement matrix. It is a nxn matrix in which rows and columns are alternatives and ranks respectively.

$$Q = [q_i \ell = \sum_{j=1}^2 G_i' \ell_j W_j] \dots\dots\dots (14) \quad \text{Where } w_i \text{ is the weight of criteria } j \text{ and } G_{i\ell j} = 1 \text{ if alternative } i \text{ is assigned rank } \ell \text{ given criterion } j. \text{ otherwise it is zero.}$$

**Step – 9 :** Formulate a mathematical model. In order to obtain final ranks of alternatives, the classical assignment problem is considered. This is a zero–one model in which decision variable ( $x_{i\ell}$ ) is one if rank  $\ell$  is assigned to alternative  $i$ , otherwise it is zero.

$$\begin{aligned} & \text{Max } \sum_{i=1}^n \sum_{\ell=1}^n q_{i\ell} x_{i\ell} \\ \text{subject to } & \sum_{i=1}^n x_{i\ell} = 1, \ell = 1,2,3 \dots\dots\dots n \\ & \sum_{\ell=1}^n x_{i\ell} = 1, i = 1,2,3 \dots\dots\dots n \end{aligned} \quad \dots\dots\dots (15)$$

$x_{i\ell}$  is binary. Solving the above model represents the final ranking of alternatives.

**Section – 3 : Numerical Example**

A company produces five products a,b,c,d,e. Demand, selling price, raw material cost and delivery cost of the products are triangular fuzzy numbers as shown in Table 1. Processing time and available capacity are shown in Table 2.

Product	Demand	Selling price (dollar)	Raw material cost (dollar)	CM (SP-RMC)	Late delivery cost (dollar)
a	(4,6,7,9,)	(3,5,7,10,)	(5,7,10,12,)	(3,4,6,9,)	9

b	(2,3,5,9,)	(5,7,9,13,)	(4,6,9,12,)	(5,6,7,10,)	4
c	(7,9,10,12,)	(6,7,9,10,)	(7,9,10,13,)	(6,7,10,13,)	2
d	(4,5,7,9,)	(5,7,12,15,)	(7,9,13,15,)	(2,4,10,13,)	1
e	(5,7,10,14,)	(4,10,13,15,)	(3,7,9,13,)	(2,3,10,14,)	8

**Table – 1** Demand, selling price, Raw material cost of each product and Late delivery cost.

Station-1	Station - 2	Station-3	Station-4	Station-5
(3,7,10,13,1)	(4,7,10,14,1)	(5,15,20,30,)	(5,10,15,20,)	(10,15,20,)
(10,15,20,30)	(5,15,20,35)	(5,10,15,25)	(10,15,20,30)	(5,15,20,25)
(5,10,20,40)	(10,15,20,35)	(10,15,25,30)	(5,10,15,25)	(10,15,25,35)
(10,15,25,30)	(10,20,30,40)	(5,10,15,25)	(10,15,25,35)	(15,20,25,40)
(5,10,15,30)	(0,0,0,0)	(0,0,0,0)	(5,10,15,20)	(0,0,0,0)

**Table – 2** Processing time of each product.

The available capacity is (300,950,2000,2500) , (250,1250,3500,4000), (150,650,1200,1800), (175,600,1000,1200) , (200,350,600,800).

**Step - :** The required capacity of each station is calculated as follows.

$$Rc_1 = \sum_{i=a}^e t_{i1}D_i = (275, 900,1900,2400)$$

$$Rc_2 = \sum_{i=a}^e t_{i2}D_i = (225, 725,1800,2200)$$

$$Rc_3 = \sum_{i=a}^e t_{i3}D_i = (200, 750,1325,1365)$$

e

$$Rc_4 = \sum_{i=a}^e t_{i4}D_i = (275, 900, 2000, 2100)$$

$$Rc_5 = \sum_{i=a}^e t_{i5}D_i = (275, 975, 1825, 2000)$$

Since  $Rc_2 > Ac_2$  ,  $Rc_3 > Ac_3$  ,  $Rc_5 > Ac_5$  so stations 2,3 and 5 are inventory level but station 1 is not. It can be easily concluded whether station 4 is an inventory level. ,  $Ac_4$  and  $Rc_4$  are compared using left and right.

$$\begin{aligned} \mu_{AC4} &= \begin{array}{l} x-175, \quad 175 < x < 600 \\ \hline 425 \\ 1000-x \quad 600 < x < 1000 \\ \hline 400 \\ 1200-x \quad 1000 < x < 1200 \\ \hline 200 \end{array} \quad \mu_{max} = \begin{array}{l} x-175 \\ \hline 1825 \end{array} \\ \mu_{RC4} &= \begin{array}{l} x-275, \quad 275 < x < 900 \\ \hline 625 \\ 2000-x \quad 900 < x < 2000 \\ \hline 1100 \\ 2100-x \quad 2000 < x < 2100 \\ \hline 100 \end{array} \quad \mu_{min} = \begin{array}{l} 2000-x \\ \hline 1825 \end{array} \end{aligned}$$

This right score of fuzzy number's are the intersecting of their non-increasing parts and  $\mu_{max}(x)$

$\mu_{Right}(Ac_4) = 0.42$  and  $\mu_{Right}(Rc_4) = 0.53$  similarly the left and total scores of  $Rc_4$  and  $Ac_4$  are calculated as



$\mu_{Left}(Ac_4) = 0.76$  and  $\mu_{Left}(Rc_1) = 0.68$ ,  $\mu_{Total}(Ac_4) = 0.32$  and  $\mu_{Total}(Rc_4) = 0.41$ . so  $Rc_4$  is greater than  $Ac_4$  and state 4 is an inventory level.

**Step 2 :** Late delivery costs are assumed crisp values, However, the algorithm would be efficient, if they were fuzzy. Decision matrices are set as follows. Note that the first column of the following matrices are calculated by dividing  $C_{mi}$  into  $t_{ij}$

$$DM_1 = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 0.32 & 2 & 12 & 5 \\ 0.36 & 2.13 & 2.7 & 2.5 \\ 0.32 & 0 & 5 & 8 \\ 0.5 & 2.22 & 8 & 8 \\ 0.42 & 2.5 & 7 & 2.4 \end{pmatrix} \quad DM_2 = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 0.4 & 6 & 8 & 5 \\ 0.42 & 2.11 & 5 & 4.2 \\ 0.28 & 0.57 & 2 & 5.3 \\ 0.42 & 2 & 9 & 2.8 \\ - & - & - & - \end{pmatrix}$$

$$DM_3 = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 0.43 & 1 & 13 & 12.3 \\ 0.43 & 2.01 & 5 & 5.4 \\ 0.1 & 0.34 & 2 & 3.9 \\ 7 & 3 & 6 & 6.8 \\ - & - & - & 3.7 \end{pmatrix} \quad DM_4 = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 0.78 & 4 & 3 & 2.3 \\ 0.35 & 1.76 & 4.6 & 7 \\ 0.54 & 2 & 4 & 4 \\ 0.43 & 1.89 & 2 & 5 \\ 0.43 & 1.7 & 7 & 8 \end{pmatrix}$$

$$DM_5 = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 0.81 & 5 & 6 & 5 \\ 0.56 & 1.76 & 8 & 0 \\ 0.9 & 0.76 & 9 & 6 \\ 0.59 & 1.76 & 0 & 7 \\ - & - & - & - \end{pmatrix}$$

**Step 3:** Ranking alternatives given the first criterion is done by applying left and right scores method. Consider the column of  $DM_1$ . It is clear that the ranks assigned to c,d, and e are 5,4,3. so the ranks assigned to 'a' and 'b' are 1 and 2 respectively. The ordinal rank matrices for  $DM_1$  are set as follows.

$$R_1 = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 4 & 1 & 2 \\ 3 & 5 & 1 \\ 2 & 2 & 4 \\ 5 & 4 & 5 \\ 1 & 3 & 3 \end{pmatrix} \quad R_2 = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 3 & 4 & 3 \\ 5 & 5 & 4 \\ 2 & 1 & 2 \\ 1 & 2 & 1 \\ 4 & 3 & 5 \end{pmatrix} \quad R_3 = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 5 & 4 & 3 \\ 4 & 3 & 4 \\ 1 & 2 & 1 \\ 2 & 1 & 5 \\ 3 & 5 & 2 \end{pmatrix}$$

**Step 4:** The agreed matrix given thought  $R_1$  is set by a

$$R_1^1 = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 5 & 1 & 3 & 5 \\ 2 & 2 & 2 & 1 \\ 1 & 5 & 5 & 4 \\ 4 & 3 & 1 & 3 \\ 3 & 4 & 4 & 2 \end{pmatrix} \quad R_2^1 = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 5 & 2 & 3 & 3 \\ 4 & 4 & 1 & 4 \\ 1 & 5 & 5 & 1 \\ 2 & 3 & 2 & 5 \\ 3 & 1 & 4 & 2 \end{pmatrix}$$

**Step 5:** Agreed matrices are converted into Borda score matrices

$$B_1 = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 3 & 4 & 4 & 4 \\ 4 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 2 & 2 & 3 \\ 2 & 3 & 3 & 2 \end{pmatrix} \quad B_2 = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 0 & 3 & 3 & 4 \\ 1 & 1 & 0 & 2 \\ 4 & 0 & 1 & 1 \\ 0 & 2 & 2 & 0 \\ 2 & 4 & 4 & 3 \end{pmatrix}$$

**Step 6:** score matrices are set by summing the values of each row.

$$Sm_1 = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 15 \\ 5 \\ 2 \\ 6 \\ 10 \end{pmatrix} \quad Sm_2 = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 10 \\ 4 \\ 6 \\ 4 \\ 13 \end{pmatrix}$$

**Step 7:** In order to get the agreement matrix the values of  $Sm_1$  and  $Sm_2$  are ranked and form the first and second columns of the agreement matrix respectively.

$$R_G = \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} \begin{pmatrix} 3 & 1.3 \\ 5 & 5 \\ 4 & 3 \\ 1 & 4 \\ 2 & 1 \end{pmatrix}$$

**Step 8:** Assume that the weight vector of criteria are (0.7,0.3). The collective weight agreement matrix is set as follows.

$$R_G = \begin{matrix} & R1 & R2 & R3 & R4 & R5 \\ \begin{matrix} a \\ b \\ c \\ d \\ e \end{matrix} & \left( \begin{matrix} \mathbf{0.6} & 0.9 & 0 & 0 & 0 \\ 0.3 & 0.3 & 0 & 0 & 0 \\ 0 & 0 & 0.2 & 0 & 0 \\ 0 & 0 & 0.3 & 0.9 & \mathbf{0.4} \\ 0 & 0 & 0 & 0.3 & 0.6 \end{matrix} \right) \end{matrix}$$

or instance  $a_{11} = 0.6$ ,  $a_{35} = 0.4$  because rank 1 is assigned to alternative ‘a’ given the first criterion.

**Step – 9 :** The assignment modal is formulated as follows.

$$\text{Max } z = 0.6 X_{a1} + 0.9X_{a2} + 0.3X_{b1} + \dots\dots\dots 0.6X_{e5}$$

$$\text{Subject to } \sum_{i=a}^e x_{i\ell} = 1, \ell = 1,2,3 \dots\dots\dots 5$$

$$\text{Subject to } \sum_{\ell=1}^5 x_{i\ell} = 1, i = a,b,c,d,e$$

The optimal solution is  $X_{a1} = X_{b2} = X_{c3} = X_{d4} = X_{e5} = 1$ . It means that ‘a’ has the highest production priority while ‘d’ has the lowest one.

**Conclusion:** The improved algorithm benefits from the advantage of reaching optimal solution. In the previous researchers all inputs of the were considered as crisp values. The assumption is not in real cases. This paper considers product mix problem as a group

decision making problem in which all inputs are fuzzy. In this paper, a new algorithm for optimizing product mix under fuzzy parameters is developed. For this method, ordering methods are used in order to make decision in a fuzzy group decision making environment.

### References :

1. Aryanezhad, M.B. and Rashidi Komijan, A (2004) “An improved Algorithm for optimizing product mix under the theory of constraints”, International journal of production Research, 42, 4221-4233.
2. Chen. S.J.and Hwang, C.L (1992) Fuzzy multiple Attribute Decision making, Springer-verlog.
3. Fredendall, L.D and Lea, B.R (117) “Improving the product mix Heuristic in the of constrains”, International journal of production research, 35, 1535-1544.
4. Goldratt.E.M (1984) The Goal, Newyork, North River press.
5. Hwang, C.L and Lin M.J (1987) , Group decision making under multiple criteria, Springer-verlag.
6. Lee.T.N. and Plenert, G(1993) ‘Optimizing theory of constraints when new product Alternatives Exist’ Production and inventory management journal, 34, 51-57.
7. Luebbe. And Finch. B (1992) ‘Theory of constraints and linear programming ‘A comparison International journal of production Research 30, 1471-1478.
8. Patterson, M.C(1992) ‘The product mix decision’ A comparison of theory of constraints and Labor – Based management Accounting’ prod and Inventory management journal 33, 80-85.
9. Rashidi Komijan. A and Sadjadi, S.J(2005) ‘optimizing product mix in a multi – bottleneck Environment proceeding of international conference on computational science and its applications, suntec, Singapore, 388-396.